



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

phorus nigricans (n), *Seriola zonata* (l, o.), *Pomatomus saltatrix* (s, b, n.), *Trachinotus falcatus* (b), *Rachycentron canadum* (b), *Poronotus triacanthus* (e), *Aphredoderus sayanus* (c), *Lepomis auritus* (c, p.), *Pomotis gibbosus* (bu, bo, w, p.), *Microp-terus dolomieu*, (w, bo, Millstone River below Rocky Hill), *M. salmoides* (bo), *Perca flavescens* (bu, bo, c, p.), *Boleosoma nigrum olmstedii* (p), *Roccus lineatus* (b, l, Barnegat), *Morone americana* (s), *Centropristis striatus* (s, b.), *Stenotomus chrysops* (s, b.), *Lagodon rhomboides* (b), *Mullus auratus* (s), *Cynoscion regalis* (l, s, b.), *Bairdiella chrysura* (s), *Leiostomus xanthurus* (s, b.), *Micropogon undulatus* (b), *Menticirrhus saxatilis* (s, e.), *Pogonias cromis* (b), *Tautogolabrus adspersus* (b), *Tautoga onitis* (l, s, b.), *Balistes carolinensis* (o), *Alutera schoepfii* (s, b, n.), *Stephanolepis hispidus* (b, s.), *Lagocephalus laevigatus* (o), *Spheroides maculatus* (o, s, b.), *Chilomycterus schoepfi* (s, b.), *Myoxocephalus octodecimspinosus* (n, m, o.), *M. aeneus* (s, b.), *Prionotus evolans strigatus* (s, b, o.), *P. carolinus* (o, s, b.), *Paralichthys dentatus* (b, s, o, l.), *Pseudopleuronectes americanus* (s), *Achirus fasciatus* (b), *Leptecheneis naucratis* (b), *Opsanus tau* (s, b, l.), *Astroscopus guttatus* (s), *Enchelyopus anguillaris* (o), *Rissola marginata* (b), *Gadus callarias* (o), *Pollachius virens* (o), *Microgadus tomcod* (s), *Urophycis regius* (s, l.), *U. chuss* (o), *U. tenuis* (o), *Merluccius bilinearis* (b, l.), *Lophius piscatorius* (o, l.), *Leptocephalus conger* (l).

HENRY W. FOWLER,
Philadelphia, Pa.

BURROWING HABITS OF THE BOX TURTLE.

Dr. Overton's note in COPEIA No. 26 on "Aquatic Habits of the Box Turtle" (*Cistudo carolina*) recalls to the writer observations on Long Island, which indicate periods of aestivation for this species.

At Yaphank box turtles can always be found in considerable numbers during hot summer weather buried to a depth of from six to ten inches in the mud bordering a pool and several springs of a cranberry bog. On short visits to the region, usually two or three days, it could not be ascertained with certainty how long a time the turtles remain buried, yet that it must be for days, and, during prolonged droughts, probably for weeks appears certain, since in many cases observed the entry holes of the turtle burrows had been almost closed through sagging, followed by drying of the mud. From such burrows the turtles could not escape, except by breaking through the crust of mud at the top.

While the burrowing habit of the box turtle is commonplace and well-known, the gathering of such numbers of this species as witnessed by the writer and Mr. Wm. T. Davis during August, 1913, on Shelter Island, no doubt, is unusual. In a short, narrow ditch, partly filled with mud and water, we counted sixty, without disturbing the turtles, and there probably were as many more packed away in the mud. New arrivals were still coming in from the surrounding wood.

GEORGE P. ENGELHARDT,
Brooklyn, N. Y.

THE BREEDING SEASON OF *BUFO* *MARINUS* (L) IN DEMERARA.

In a recent number of *COPEIA* (February 24, 1916) Mr. Austin H. Clark says of the breeding season of *Bufo aqua* (*Bufo marinus* L.) "appears to breed about the commencement of the rainy season, somewhere in November or thereabouts," crediting this statement to Mr. Gilbert E. Bodkin, Government Economic Biologist of British Guiana.

The writer has no data on the habits of the species in Demerara, except for the months of July, August and September, but it should be recorded that